

# EXPONENTIAL MAP AND $L_\infty$ ALGEBRA ASSOCIATED TO A LIE PAIR

CAMILLE LAURENT-GENGOUX, MATHIEU STIÉNON, AND PING XU

Dédié à la mémoire de Paulette Libermann.

**ABSTRACT.** In this note, we unveil homotopy-rich algebraic structures generated by the Atiyah classes relative to a Lie pair  $(L, A)$  of algebroids. In particular, we prove that the quotient  $L/A$  of such a pair admits an essentially canonical homotopy module structure over the Lie algebroid  $A$ , which we call Kapranov module.

*Camille Laurent-Gengoux, Mathieu Stiénon, Ping Xu, C. R. Acad. Sci. Paris, Ser. I 350 (2012) 817-821.*

## 1. KAPRANOV MODULES

Let  $A$  be a Lie algebroid (either real or complex) over a manifold  $M$  with anchor  $\rho$ . By an  $A$ -module, we mean a module of the corresponding Lie-Rinehart algebra  $\Gamma(A)$  over the associative algebra  $C^\infty(M)$ . An  $A$ -connection on a smooth vector bundle  $E$  over  $M$  is a bilinear map  $\nabla : \Gamma(A) \otimes \Gamma(E) \rightarrow \Gamma(E)$  satisfying  $\nabla_{fa}e = f\nabla_ae$  and  $\nabla_a(fe) = (\rho(a)f)e + f\nabla_ae$ , for all  $a \in \Gamma(A)$ ,  $e \in \Gamma(E)$ , and  $f \in C^\infty(M)$ . A vector bundle  $E$  endowed with a *flat*  $A$ -connection (also known as an infinitesimal  $A$ -action) is an  $A$ -module; more precisely, its space of smooth sections  $\Gamma(E)$  is one. **Atiyah class.** Given a Lie pair  $(L, A)$  of algebroids, *i.e.* a Lie algebroid  $L$  with a Lie subalgebroid  $A$ , the Atiyah class  $\alpha_E$  of an  $A$ -module  $E$  relative to the pair  $(L, A)$  is defined as the obstruction to the existence of an  *$A$ -compatible*  $L$ -connection on  $E$ . An  $L$ -connection  $\nabla$  is  $A$ -compatible if its restriction to  $\Gamma(A) \otimes \Gamma(E)$  is the given infinitesimal  $A$ -action on  $E$  and  $\nabla_a\nabla_l - \nabla_l\nabla_a = \nabla_{[a,l]}$  for all  $a \in \Gamma(A)$  and  $l \in \Gamma(L)$ . This fairly recently defined class (see [3]) has as double origin, which it generalizes, the Atiyah class of holomorphic vector bundles and the Molino class of foliations. The quotient  $L/A$  of the Lie pair  $(L, A)$  is an  $A$ -module [3]. Its Atiyah class  $\alpha_{L/A}$  can be described as follows. Choose an  $L$ -connection  $\nabla$  on  $L/A$  extending the  $A$ -action. Its curvature is the vector bundle map  $R^\nabla : \wedge^2 L \rightarrow \text{End}(E)$  defined by  $R^\nabla(l_1, l_2) = \nabla_{l_1}\nabla_{l_2} - \nabla_{l_2}\nabla_{l_1} - \nabla_{[l_1, l_2]}$ , for all  $l_1, l_2 \in \Gamma(L)$ . Since  $L/A$  is an  $A$ -module,  $R^\nabla$  vanishes on  $\wedge^2 A$  and, therefore, determines a section  $R_{L/A}^\nabla$  of  $A^* \otimes (L/A)^* \otimes \text{End}(L/A)$ . It was proved in [3] that  $R_{L/A}^\nabla$  is a 1-cocycle for the Lie algebroid  $A$  with values in the  $A$ -module  $(L/A)^* \otimes \text{End}(L/A)$  and that its cohomology class  $\alpha_{L/A} \in H^1(A; (L/A)^* \otimes \text{End}(L/A))$  is independent of the choice of the connection.

---

Research partially supported by the National Science Foundation [DMS-1101827].

**Kapranov modules over a Lie algebroid.** Let  $M$  be a smooth manifold, and let  $R$  be the algebra of smooth functions on  $M$  valued in  $\mathbb{R}$  (or  $\mathbb{C}$ ). Let  $A$  be a Lie algebroid over  $M$ . The Chevalley-Eilenberg differential  $d_A$  and the exterior product make  $\Gamma(\wedge^\bullet A^*)$  into a differential graded commutative  $R$ -algebra.

Now let  $E$  be a smooth vector bundle over  $M$ . Deconcatenation defines an  $R$ -coalgebra structure on  $\Gamma(S^\bullet E)$ . The comultiplication  $\Delta : \Gamma(S^\bullet E) \rightarrow \Gamma(S^\bullet E) \otimes_R \Gamma(S^\bullet E)$  is given by

$$\Delta(e_1 \odot e_2 \odot \cdots \odot e_n) = \sum_{p+q=n} \sum_{\sigma \in \mathfrak{S}_p^q} (e_{\sigma(1)} \odot \cdots \odot e_{\sigma(p)}) \otimes (e_{\sigma(p+1)} \odot \cdots \odot e_{\sigma(n)}),$$

for any  $e_1, \dots, e_n \in \Gamma(E)$ . Let  $\mathfrak{e}$  denote the ideal of  $\Gamma(S^\bullet(E^*))$  generated by  $\Gamma(E^*)$ . The algebra  $\text{Hom}_R(\Gamma(S^\bullet E), R)$  dual to the coalgebra  $\Gamma(S^\bullet E)$  is the  $\mathfrak{e}$ -adic completion of  $\Gamma(S^\bullet(E^*))$ . It will be denoted by  $\Gamma(\hat{S}^\bullet(E^*))$ . Equivalently, one can think of the completion  $\hat{S}^\bullet(E^*)$  of  $S^\bullet(E^*)$  as a bundle of algebras over  $M$ . Note that  $\Gamma(\wedge^\bullet A^*)$  is an  $R$ -subalgebra of  $\Gamma(\wedge^\bullet A^* \otimes \hat{S}^\bullet E^*)$ .

Recall that an  $L_\infty[1]$  algebra is a  $\mathbb{Z}$ -graded vector space  $V = \bigoplus_{n \in \mathbb{Z}} V_n$  endowed with a sequence  $(\lambda_k)_{k=1}^\infty$  of symmetric multilinear maps  $\lambda_k : \otimes^k V \rightarrow V$  of degree 1 satisfying the generalized Jacobi identity

$$\sum_{k=1}^n \sum_{\sigma \in \mathfrak{S}_k^{n-k}} \varepsilon(\sigma; v_1, \dots, v_n) \lambda_{1+n-k}(\lambda_k(v_{\sigma(1)}, \dots, v_{\sigma(k)}), v_{\sigma(k+1)}, \dots, v_{\sigma(n)}) = 0$$

for each  $n \in \mathbb{N}$  and for any homogeneous vectors  $v_1, v_2, \dots, v_n \in V$ . Here  $\mathfrak{S}_p^q$  denotes the set of  $(p, q)$ -shuffles<sup>1</sup> and  $\varepsilon(\sigma; v_1, \dots, v_n)$  the Koszul sign<sup>2</sup> of the permutation  $\sigma$  of the (homogeneous) vectors  $v_1, v_2, \dots, v_n$ .

**Definition 1.1.** A Kapranov module over a Lie algebroid  $A \rightarrow M$  is a vector bundle  $E \rightarrow M$  together with an  $L_\infty[1]$  algebra structure on  $\Gamma(\wedge^\bullet A^* \otimes E)$  defined by a sequence  $(\lambda_k)_{k \in \mathbb{N}}$  of multibrackets (called Kapranov multibrackets) such that (1) the unary bracket  $\lambda_1 : \Gamma(\wedge^\bullet A^* \otimes E) \rightarrow \Gamma(\wedge^{\bullet+1} A^* \otimes E)$  is the Chevalley-Eilenberg differential associated to an infinitesimal  $A$ -action on  $E$ , and (2) all multibrackets  $\lambda_k : \otimes^k \Gamma(\wedge^\bullet A^* \otimes E) \rightarrow \Gamma(\wedge^\bullet A^* \otimes E)[1]$  with  $k \geq 2$  are  $\Gamma(\wedge^\bullet A^*)$ -multilinear.

**Proposition 1.1.** Let  $A$  be a Lie algebroid over a smooth manifold  $M$  and let  $E$  be a smooth vector bundle over  $M$ . Each of the following four data is equivalent to a Kapranov  $A$ -module structure on  $E$ .

- (1) A degree 1 derivation  $D$  of the graded algebra  $\Gamma(\wedge^\bullet A^* \otimes \hat{S}(E^*))$ , which preserves the filtration  $\Gamma(\wedge^\bullet A^* \otimes \hat{S}^{\geq n}(E^*))$ , satisfies  $D^2 = 0$ , and whose restriction to  $\Gamma(\wedge^\bullet A^*)$  is the Chevalley-Eilenberg differential of the Lie algebroid  $A$ . (Here, by convention, all elements of  $\hat{S}(E^*)$  have degree 0.)
- (2) An infinitesimal action of  $A$  on  $\hat{S}(E^*)$  by derivations which preserve the decreasing filtration  $\hat{S}^{\geq n}(E^*)$ .
- (3) An infinitesimal action of  $A$  on  $S(E)$  by coderivations which preserve  $S^{\geq 1}(E)$  and the increasing filtration  $S^{\leq n}(E)$ .

1. A  $(p, q)$ -shuffle is a permutation  $\sigma$  of the set  $\{1, 2, \dots, p+q\}$  such that  $\sigma(1) \leq \sigma(2) \leq \dots \leq \sigma(p)$  and  $\sigma(p+1) \leq \sigma(p+2) \leq \dots \leq \sigma(p+q)$ .

2. The Koszul sign of a permutation  $\sigma$  of the (homogeneous) vectors  $v_1, v_2, \dots, v_n$  is determined by the relation  $v_{\sigma(1)} \odot v_{\sigma(2)} \odot \cdots \odot v_{\sigma(n)} = \varepsilon(\sigma; v_1, \dots, v_n) v_1 \odot v_2 \odot \cdots \odot v_n$ .

- (4) An infinitesimal action of  $A$  on  $E$  together with a sequence of morphisms of vector bundles  $\mathbf{R}_k : S^k(E) \rightarrow A^* \otimes E$  ( $k \geq 2$ ) whose sum

$$\mathbf{R} = \sum_{k=2}^{\infty} \mathbf{R}_k \in \Gamma(A^* \otimes \hat{S}(E^*) \otimes E)$$

is a solution of the Maurer-Cartan equation  $d_A \mathbf{R} + \frac{1}{2} [\mathbf{R}, \mathbf{R}] = 0$ . (Here, we consider  $\Gamma(\hat{S}(E^*) \otimes E)$  as the space of formal vertical vector fields on  $E$  along the zero section and derive a natural Lie bracket on the graded vector space  $\Gamma(\wedge^\bullet A^* \otimes \hat{S}(E^*) \otimes E)$ .)

Characterizations (1) and (4) are related by the identity  $D = d_A^{\hat{S}(E^*)} + \mathbf{R}$ , where  $d_A^{\hat{S}(E^*)}$  denotes the Chevalley-Eilenberg differential associated to the infinitesimal  $A$ -action on  $E$ , and  $\mathbf{R}$  denotes its own action on  $\Gamma(\wedge^\bullet A^* \otimes \hat{S}(E^*))$  by contraction. On the other hand, for any  $k \geq 2$ , the  $k$ -th Kapranov multibracket  $\lambda_k$  is related to the  $k$ -th component  $\mathbf{R}_k \in \Gamma(A^* \otimes S^k E^* \otimes E)$  of the Maurer-Cartan element  $\mathbf{R}$  through the equation

$$\lambda_k(\xi_1 \otimes e_1, \dots, \xi_k \otimes e_k) = (-1)^{|\xi_1| + \dots + |\xi_k|} \xi_1 \wedge \dots \wedge \xi_k \wedge \mathbf{R}_k(e_1, \dots, e_k),$$

which is valid for any  $e_1, \dots, e_k \in \Gamma(E)$  and any homogeneous elements  $\xi_1, \dots, \xi_k$  of  $\Gamma(\wedge^\bullet A^*)$ .

The algebraic structure described in the above proposition is related to Costello's  $L_\infty$  algebras over the differential graded algebra  $(\Gamma(\wedge^\bullet A^*), d_A)$  [4], and to Yu's  $L_\infty$  algebroids [9].

Two Kapranov  $A$ -modules  $E_1$  and  $E_2$  over  $M$  are isomorphic if there exists an isomorphism  $\Phi : S(E_1) \rightarrow S(E_2)$  of bundles of coalgebras over  $M$ , which intertwines the infinitesimal  $A$ -actions.

## 2. EXPONENTIAL MAP AND POINCARÉ-BIRKHOFF-WITT ISOMORPHISM

Assume  $\mathcal{A}$  is a Lie subgroupoid of a Lie groupoid  $\mathcal{L}$  (over the same unit space), and let  $A$  and  $L$  denote the corresponding Lie algebroids. The source map  $s : \mathcal{L} \rightarrow M$  factors through the quotient of the action of  $\mathcal{A}$  on  $\mathcal{L}$  by multiplication from the right. Therefore, it induces a surjective submersion  $s : \mathcal{L}/\mathcal{A} \rightarrow M$ . Note that the zero section  $0 : M \rightarrow L/A$  and the unit section  $1 : M \rightarrow \mathcal{L}/\mathcal{A}$  are both embeddings of  $M$ .

**Proposition 2.1.** *Each choice of a splitting of the short exact sequence of vector bundles  $0 \rightarrow A \rightarrow L \rightarrow L/A \rightarrow 0$  and of an  $L$ -connection  $\nabla$  on  $L/A$  extending the  $A$ -action determines an exponential map, i.e. a fiber bundle map*

$$\exp^\nabla : L/A \rightarrow \mathcal{L}/\mathcal{A},$$

*which identifies the zero section of  $L/A$  to the unit section of  $\mathcal{L}/\mathcal{A}$ , whose differential along the zero section of  $L/A$  is the canonical isomorphism between  $L/A$  and the tangent bundle to the  $s$ -foliation of  $\mathcal{L}/\mathcal{A}$  along the unit section, and which is locally diffeomorphic around  $M$ .*

Let  $\mathcal{N}(L/A)$  denote the space of all functions on  $L/A$  which, together with their derivatives of all degrees in the direction of the  $\pi$ -fibers, vanish along the zero section. The space of  $\pi$ -fiberwise differential operators on  $L/A$  along the zero

section is canonically identified to the symmetric  $R$ -algebra  $\Gamma(S(L/A))$ . Therefore, we have the short exact sequence of  $R$ -algebras

$$(1) \quad 0 \rightarrow \mathcal{N}(L/A) \rightarrow C^\infty(L/A) \rightarrow \text{Hom}_R(\Gamma(S(L/A)), R) \rightarrow 0.$$

Likewise, let  $\mathcal{N}(\mathcal{L}/\mathcal{A})$  denote the space of all functions on  $\mathcal{L}/\mathcal{A}$  which, together with their derivatives of all degrees in the direction of the  $s$ -fibers, vanish along the unit section. The space of  $s$ -fiberwise differential operators on  $\mathcal{L}/\mathcal{A}$  along the unit section is canonically identified to the quotient of the enveloping algebra  $\mathcal{U}(L)$  by the left ideal generated by  $\Gamma(A)$ . Therefore, we have the short exact sequence of  $R$ -algebras

$$(2) \quad 0 \rightarrow \mathcal{N}(\mathcal{L}/\mathcal{A}) \rightarrow C^\infty(\mathcal{L}/\mathcal{A}) \rightarrow \text{Hom}_R\left(\frac{\mathcal{U}(L)}{\mathcal{U}(L)\Gamma(A)}, R\right) \rightarrow 0.$$

Since the exponential (or more precisely its dual) maps  $\mathcal{N}(\mathcal{L}/\mathcal{A})$  to  $\mathcal{N}(L/A)$ , it induces an isomorphism of  $R$ -modules from  $\text{Hom}_R(\Gamma(S(L/A)), R)$  to  $\text{Hom}_R\left(\frac{\mathcal{U}(L)}{\mathcal{U}(L)\Gamma(A)}, R\right)$ .

**Proposition 2.2.** *Each choice of a splitting of the short exact sequence of vector bundles  $0 \rightarrow A \rightarrow L \rightarrow L/A \rightarrow 0$  and of an  $L$ -connection  $\nabla$  on  $L/A$  extending the  $A$ -action determines an isomorphism of filtered  $R$ -modules*

$$\text{PBW} : \Gamma(S(L/A)) \rightarrow \frac{\mathcal{U}(L)}{\mathcal{U}(L)\Gamma(A)}$$

*called Poincaré-Birkhoff-Witt map.*

**Remark 2.1.** *In case  $L = A \bowtie B$  is the Lie algebroid sum of a matched pair of Lie algebroids  $(A, B)$ , the  $L$ -connection  $\nabla$  on  $L/A \cong B$  extending the  $A$ -action determines a  $B$ -connection on  $B$ , the coalgebras  $\frac{\mathcal{U}(L)}{\mathcal{U}(L)\Gamma(A)}$  and  $\mathcal{U}(B)$  are isomorphic, and the corresponding Poincaré-Birkhoff-Witt map  $\text{PBW} : \Gamma(S(B)) \rightarrow \mathcal{U}(B)$  is standard (see [7] for instance).*

**Proposition 2.3.** *The Poincaré-Birkhoff-Witt map associated to a splitting  $j : L/A \rightarrow L$  of the short exact sequence of vector bundles  $0 \rightarrow A \rightarrow L \rightarrow L/A \rightarrow 0$  and an  $L$ -connection  $\nabla$  on  $L/A$  satisfies  $\text{PBW}(1) = 1$  and, for all  $b \in \Gamma(L/A)$  and  $n \in \mathbb{N}$ ,  $\text{PBW}(b) = j(b)$  and  $\text{PBW}(b^{n+1}) = j(b) \cdot \text{PBW}(b^n) - \text{PBW}(\nabla_{j(b)}(b^n))$ , where  $b^k$  stands for the symmetric product  $b \odot b \odot \cdots \odot b$  of  $k$  copies of  $b$ .*

**Remark 2.2.** *Although the construction of the Poincaré-Birkhoff-Witt map outlined above presupposes that  $L$  and  $A$  are integrable real Lie algebroids, PBW can be defined for any real (resp. complex) Lie pair provided one works with local (resp. formal) groupoids.*

Recall that  $L/A$  is an  $A$ -module. The representation of  $A$  on  $L/A$ , an obvious generalization of the Bott connection, induces an infinitesimal action of  $A$  on the coalgebra  $\Gamma(S(L/A))$  by coderivation. Besides, multiplication by elements of  $\Gamma(A)$  from the left in the coalgebra  $\mathcal{U}(L)$  induces an infinitesimal action of  $A$  on the coalgebra  $\frac{\mathcal{U}(L)}{\mathcal{U}(L)\Gamma(A)}$  by coderivation.

The infinitesimal actions of  $A$  on  $L/A$  and  $\mathcal{L}/\mathcal{A}$  induce infinitesimal actions of  $A$  by derivations on the algebras of functions  $C^\infty(L/A)$  and  $C^\infty(\mathcal{L}/\mathcal{A})$  and, consequently, on the algebras of infinite jets  $\text{Hom}_R(\Gamma(S(L/A)), R)$  and  $\text{Hom}_R\left(\frac{\mathcal{U}(L)}{\mathcal{U}(L)\Gamma(A)}, R\right)$ .

**Proposition 2.4.** (1) The space  $\text{Hom}_R(\frac{\mathcal{U}(L)}{\mathcal{U}(L)\Gamma(A)}, R)$  of infinite  $s$ -fiberwise jets along  $M$  of functions on  $\mathcal{L}/\mathcal{A}$  is an associative algebra on which the Lie algebroid  $A$  acts infinitesimally by derivations. (2) The dual of the exponential map  $\text{PBW}^* : \text{Hom}_R(\frac{\mathcal{U}(L)}{\mathcal{U}(L)\Gamma(A)}, R) \rightarrow \text{Hom}_R(\Gamma(S(L/A)), R)$  is an isomorphism of associative algebras, which may or may not intertwine the infinitesimal  $A$ -actions.

### 3. $L_\infty[1]$ ALGEBRA ASSOCIATED TO A LIE PAIR

Our main result is the following

**Theorem 3.1.** If  $(L, A)$  is a Lie pair, i.e. a Lie algebroid  $L$  together with a Lie subalgebroid  $A$ , then  $L/A$  admits a Kapranov module structure, canonical up to isomorphism, over the Lie algebroid  $A$ , whose  $\mathbf{R}_2 \in \Gamma(A^* \otimes S^2(L/A)^* \otimes L/A)$  (see Proposition 1.1) is a 1-cocycle representative of the Atiyah class of  $L/A$  relative to the pair  $(L, A)$ .

Moreover, when  $L = A \bowtie B$  is the Lie algebroid sum of a matched pair  $(A, B)$  of Lie algebroids and there exists a torsion free flat  $B$ -connection  $\nabla$  on  $B$ , the components of the Maurer-Cartan element  $\mathbf{R}$  satisfy the recursive formula  $\mathbf{R}_{k+1} = \partial^\nabla \mathbf{R}_k$ , where  $\partial^\nabla$  denotes the covariant differential associated to the connection.

**Sketch of proof** Choose a splitting of the short exact sequence of vector bundles  $0 \rightarrow A \rightarrow L \rightarrow L/A \rightarrow 0$  and an  $L$ -connection  $\nabla$  on  $L/A$  extending the  $A$ -action. Identify  $\Gamma(\hat{S}(L/A)^*)$  to  $\text{Hom}_R(\frac{\mathcal{U}(L)}{\mathcal{U}(L)\Gamma(A)}, R)$  via the PBW map and pull back the infinitesimal  $A$ -action of the latter to the former. According to Proposition 1.1, the resulting  $A$ -action on  $\Gamma(\hat{S}(L/A)^*)$  by derivations determines a Kapranov  $A$ -module structure on  $L/A$ . Making use of Proposition 2.3, one can check directly that  $\mathbf{R}_2$  is a 1-cocycle representative of the Atiyah class  $\alpha_{L/A}$ .  $\square$

As immediate consequences, we recover the following results of [3].

**Corollary 3.2.** Given a Lie algebroid pair  $(L, A)$ , let  $\mathcal{U}(A)$  denote the universal enveloping algebra of the Lie algebroid  $A$  and let  $\mathcal{A}$  denote the category of  $\mathcal{U}(A)$ -modules. The Atiyah class of the quotient  $L/A$  makes  $L/A[-1]$  into a Lie algebra object in the derived category  $D^b(\mathcal{A})$ .

**Corollary 3.3.** Let  $(L, A)$  be a Lie pair and let  $\mathcal{C}$  be a bundle (of finite or infinite rank) of associative commutative algebras on which  $A$  acts by derivations. There exists an  $L_\infty[1]$  algebra structure on  $\Gamma(\wedge^\bullet A^* \otimes L/A \otimes \mathcal{C})$ , canonical up to  $L_\infty$  isomorphism. Moreover,  $H^{\bullet-1}(A; L/A \otimes \mathcal{C})$  is a graded Lie algebra whose Lie bracket only depends on the Atiyah class of  $L/A$ .

For  $\mathcal{C} = \mathbb{C}$ , the Lie bracket on the cohomology  $H^{\bullet-1}(A, L/A)$  happens to be trivial.

### 4. AN EXAMPLE DUE TO KAPRANOV

Let  $X$  be a Kähler manifold with real analytic metric. Recall that the eigenbundles  $T_X^{0,1}$  and  $T_X^{1,0}$  of the complex structure  $J : T_X \rightarrow T_X$  ( $J^2 = -\text{id}$ ) form a matched pair of Lie algebroids [6]. Fix a point  $x \in X$ . The exponential map  $\exp_x^{\text{LC}} : T_x X \rightarrow X$  defined using the geodesics of the Levi-Civita connection  $\nabla^{\text{LC}}$  originating from the point  $x$  needs not be holomorphic.

However, Calabi constructed a holomorphic exponential map  $\exp_x^{\text{hol}} : T_x X \rightarrow X$  as follows [2] (see also [1]). First, extend the Levi-Civita connection  $\mathbb{C}$ -linearly to a

$T_X \otimes \mathbb{C}$ -connection  $\nabla^{\mathbb{C}}$  on  $T_X \otimes \mathbb{C}$ . Since  $X$  is Kähler,  $\nabla^{\text{LC}} J = 0$  and  $\nabla^{\mathbb{C}}$  restricts to a  $T_X \otimes \mathbb{C}$ -connection on  $T_X^{1,0}$ . It is easy to check that the induced  $T_X^{0,1}$ -connection on  $T_X^{1,0}$  is the canonical infinitesimal  $T_X^{0,1}$ -action on  $T_X^{1,0}$  — a section of  $T_X^{1,0}$  is  $T_X^{0,1}$ -horizontal iff it is holomorphic — while the induced  $T_X^{1,0}$ -connection  $\nabla^{1,0}$  on  $T_X^{1,0}$  is flat and torsion free. Now let  $X'$  denote the manifold  $X$  and let  $X''$  denote  $X$  with the opposite complex structure  $-J$ . The image of the diagonal embedding  $X \hookrightarrow X' \times X''$  is totally real so  $X' \times X''$  can be seen as a complexification of  $X$ . The restriction of  $T_{X' \times X''}$  (resp. its subbundle  $T_{X'} \times X''$ ) along the diagonal  $X$  is precisely the complexified tangent bundle  $T_X \otimes \mathbb{C}$  (resp. its subbundle  $T_X^{1,0}$ ). (See [8] for a discussion on integration of complex Lie algebroids.) The analytic continuation of the  $T_X^{1,0}$ -connection  $\nabla^{1,0}$  on  $T_X^{1,0}$  in a neighborhood of the diagonal is a holomorphic  $T_{X'} \times X''$ -connection on the Lie algebroid  $T_{X'} \times X''$ , whose exponential map  $\exp_x^{\text{hol}}$  at a diagonal point  $(x, x)$  takes  $T_x X' \times \{x\}$  (which is  $(T_X^{1,0})_x$  or  $T_x X$ ) into  $X' \times \{x\}$  (which is  $X$ ).

Consider the Lie pair  $(L = T_{X' \times X''}, A = X' \times T_{X''})$ , the corresponding Lie groupoids  $\mathcal{L} = (X' \times X'') \times (X' \times X'')$  and  $\mathcal{A} = X' \times (X'' \times X'')$ , and the associated quotients  $L/A = T_{X'} \times X''$  and  $\mathcal{L}/\mathcal{A} = (X' \times X') \times X''$ . Calabi's holomorphic exponential map  $\exp^{\text{hol}}$  is indeed the restriction along the diagonal of the exponential map  $\exp^{\nabla^{1,0}} : L/A \rightarrow \mathcal{L}/\mathcal{A}$  associated to the  $T_{X'} \times X''$ -connection  $\nabla^{1,0}$  on the Lie algebroid  $T_{X'} \times X''$  as described in Proposition 2.1.

Taking the infinite jet of  $\exp^{\text{hol}}$ , we obtain, as in Proposition 2.2, a Poincaré-Birkhoff-Witt map  $\text{PBW}^{\text{hol}} : \Gamma(S(T_X^{1,0})) \rightarrow \mathcal{U}(T_X^{1,0})$ . Then, pulling back the infinitesimal  $T_X^{0,1}$ -action on  $\mathcal{U}(T_X^{1,0})$  to an infinitesimal  $T_X^{0,1}$ -action by coderivations on  $\Gamma(S(T_X^{1,0}))$ , we obtain, as in Theorem 3.1, a Kapranov  $T_X^{0,1}$ -module structure on  $T_X^{1,0}$ . In this context, the tensors  $R_n \in \Omega^{0,1}(\text{Hom}(S^n T_X^{1,0}, T_X^{1,0}))$  are the curvature  $R_2 \in \Omega^{1,1}(\text{End}(T_X^{1,0}))$  and its higher covariant derivatives. Hence we recover the following result of Kapranov:

**Theorem 4.1** ([5]). *The Dolbeault complex  $\Omega^{0,\bullet}(T_X^{1,0})$  of a Kähler manifold is an  $L_\infty[1]$  algebra. For  $n \geq 2$ , the  $n$ -th multibracket  $\lambda_n$  is the composition*

$$\Omega^{0,j_1}(T_X^{1,0}) \otimes \dots \otimes \Omega^{0,j_n}(T_X^{1,0}) \rightarrow \Omega^{0,j_1+\dots+j_n}(\otimes^n T_X^{1,0}) \rightarrow \Omega^{0,j_1+\dots+j_n+1}(T_X^{1,0})$$

*of the wedge product with the map induced by  $R_n \in \Omega^{0,1}(\text{Hom}(\otimes^n T_X^{1,0}, T_X^{1,0}))$ , while  $\lambda_1$  is the Dolbeault operator  $\bar{\partial} : \Omega^{0,j}(T_X^{1,0}) \rightarrow \Omega^{0,j+1}(T_X^{1,0})$ .*

#### ACKNOWLEDGEMENTS

Camille Laurent-Gengoux likes to express his gratitude to Penn State for its hospitality.

#### REFERENCES

1. M. Bershadsky, S. Cecotti, H. Ooguri, and C. Vafa, *Kodaira-Spencer theory of gravity and exact results for quantum string amplitudes*, Comm. Math. Phys. **165** (1994), no. 2, 311–427. MR 1301851 (95f:32029)
2. Eugenio Calabi, *Isometric imbedding of complex manifolds*, Ann. of Math. (2) **58** (1953), 1–23. MR 0057000 (15,160c)
3. Zhuo Chen, Mathieu Stiénon, and Ping Xu, *From Atiyah Classes to Homotopy Leibniz Algebras*, arXiv:1204.1075 (2012).
4. Kevin J Costello, *A geometric construction of the witten genus, II*, arXiv:1112.0816 (2011).

5. M. Kapranov, *Rozansky-Witten invariants via Atiyah classes*, Compositio Math. **115** (1999), no. 1, 71–113. MR 1671737 (2000h:57056)
6. Camille Laurent-Gengoux, Mathieu Stiénon, and Ping Xu, *Holomorphic Poisson manifolds and holomorphic Lie algebroids*, Int. Math. Res. Not. IMRN (2008), Art. ID rnn 088, 46. MR 2439547 (2009i:53082)
7. Victor Nistor, Alan Weinstein, and Ping Xu, *Pseudodifferential operators on differential groupoids*, Pacific J. Math. **189** (1999), no. 1, 117–152. MR 1687747 (2000c:58036)
8. Alan Weinstein, *The integration problem for complex Lie algebroids*, From geometry to quantum mechanics, Progr. Math., vol. 252, Birkhäuser Boston, Boston, MA, 2007, pp. 93–109. MR 2285039 (2008e:58025)
9. Shilin Yu, *Dolbeault dga of formal neighborhoods and  $L_\infty$  algebroids*, Ph.D. thesis, Penn State University, State College, PA, 2013.

DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ DE LORRAINE  
*E-mail address:* `camille.laurent-gengoux@univ-lorraine.fr`

DEPARTMENT OF MATHEMATICS, PENN STATE UNIVERSITY  
*E-mail address:* `stienon@math.psu.edu`

DEPARTMENT OF MATHEMATICS, PENN STATE UNIVERSITY  
*E-mail address:* `ping@math.psu.edu`